Chasing Serendipity: Tackling Transient Sources with Neutrino Telescopes

arxiv:2505.24666/arxiv:2505.24652

Gustavo F. S. Alves

In collaboration with:
Lua F. T. Airoldi
Yuber F. Perez-Gonzalez
Gabriel M. Salla
Renata Zukanovich Funchal

First question of the audience

First question of the audience
What is serendipity?

First question of the audience
What is serendipity?

“The occurrence and development of events by chance in a happy or beneficial way.”

“An aptitude for making desirable discoveries by accident.”

Motivation: Multi Messenger astronomy

Photons + Neutrinos

Motivation: ~~Multi~~ Messenger astronomy

~~Photons +~~ Neutrinos

Can neutrinos be the only candidate for observing a source?

Probing the universe through photons



Probing the universe through photons



Key concepts:

Probing the universe through photons



Key concepts:

- Field of view

Probing the universe through photons



Key concepts:

- Field of view

- Coordinates

Probing the universe through photons



Key concepts:

- Field of view

- Coordinates

- Interactions

The field of view

The field of view

Portion of the sky the experiment monitors

The (instantaneous) field of view

A source may be within the experiment field of view



The (instantaneous) field of view

As Earth rotates this may change

Their instantaneous field of view



Their instantaneous field of view



“latitude”

“longitude”

Their instantaneous blind spots



Their instantaneous blind spots



We will comment on the effect of varying time soon.

Their instantaneous field of view



The KM3-230213A event

KM3NeT, Nature 638, 376-382

“The signal of the most energetic elementary particle ever observed.”

KM3NeT video archive

The KM3-230213A event

$$E_\mu = 120 \text{ PeV}$$
$$\theta = 0.6^\circ$$



Serendipity in practice

Our work started before the KM3NeT paper, but after their result we have a highly motivated example to study.

Serendipity in practice

Our work started before the KM3NeT paper, but after their result we have a highly motivated example to study.

****However, everything we have discussed so far is always true**

Summary, so far.



Summary, so far.



We showed that there are blind regions in the instantaneous field of view of gamma ray experiments

Summary, so far.



We showed that there are blind regions in the instantaneous field of view of gamma ray experiments

But why instantaneous?

Ultra high energy unknown in particle physics

Ultra high energy unknown in particle physics

Clash of Titans

Ultra high energy unknown in particle physics

Clash of Titans



Ultra high energy unknown in particle physics

Clash of Titans





10% of strings!!

Ultra high energy unknown in particle physics

Origin?

Schwemberger, arxiv:2502.04508
Lu, arxiv:2509.12628

Ultra high energy unknown in particle physics

Diffuse ~ $2.5 - 3.5\ \sigma$

Origin?

Schwemberger, arxiv:2502.04508
Lu, arxiv:2509.12628

Ultra high energy unknown in particle physics

Origin?

Diffuse ~ $2.5 - 3.5\ \sigma$

Cosmogenic ~ $3.1 - 3.6\ \sigma$

Schwemberger, arxiv:2502.04508
Lu, arxiv:2509.12628

Ultra high energy unknown in particle physics

Origin?

Diffuse ~ $2.5 - 3.5\ \sigma$

Cosmogenic ~ $3.1 - 3.6\ \sigma$

Transient ~ $2\ \sigma$

Schwemberger, arxiv:2502.04508
Lu, arxiv:2509.12628

Ultra high energy unknown in particle physics

Origin?

Diffuse ~ $2.5 - 3.5\ \sigma$

Cosmogenic ~ $3.1 - 3.6\ \sigma$

Transient ~ $2\ \sigma$

A transient burst helps in recoinciling the (non) observation.

Schwemberger, arxiv:2502.04508
Lu, arxiv:2509.12628

Ultra high energy unknown in particle physics

Origin?

Diffuse ~ $2.5 - 3.5\ \sigma$

Cosmogenic ~ $3.1 - 3.6\ \sigma$

Transient ~ $2\ \sigma$

A transient burst helps in recoinciling the (non) observation.

***For fast transient signals, instantaneous field of view is what matters!**

Schwemberger, arxiv:2502.04508
Lu, arxiv:2509.12628

One example: Primordial Black Holes



The PBH evaporates



The PBH evaporates



The PBH evaporates



The PBH evaporates



Secondary vs primary

Temperature is low when it is heavy





Temperature is high when it is light

$$T \sim 1 \text{ TeV} \left(\frac{10^{10} \text{ g}}{\text{M}} \right)$$



The (neutrino) burst



BlackHawk, arxiv:1905.04268/2108.02737

The “instantaneous field of view” of neutrino telescopes



We start with IceCube

IceCube, arxiv:2101.09836





The effective area rotation

$$(\delta, \mathrm{RA})$$

The effective area rotation

$$(\delta, \mathrm{RA})$$



The effective area rotation

$$(\delta, \mathrm{RA})$$



The effective area rotation

$$(\delta, \mathrm{RA})$$



$$\theta_z$$

The effective area rotation



(δ, RA)



θ_z

Plenum, arxiv:2503.07549/2107.13534

KM3NeT effective area





10 hours later…





Number of events

Number of events

$$N_{\text{evts}} = \frac{1}{4\pi d^2} \int dE\ A(E, \delta, \text{RA}) \int_0^{\tau_B} dt \frac{dN}{dt dE}\Big|_{\text{Earth}}$$

Number of events

$$N_{\text{evts}} = \frac{1}{4\pi d^2} \int dE\ A(E, \delta, \text{RA}) \int_0^{\tau_B} dt \frac{dN}{dt dE}\Big|_{\text{Earth}}$$

What is the relevant τ_B ?

The high energy tail

$$T \simeq 7.8 \times 10^3 \times \left(\frac{1\ \mathrm{s}}{\tau_B} \right)^{1/3} \mathrm{GeV}$$

The high energy tail

$$T \simeq 7.8 \times 10^3 \times \left(\frac{1\ \mathrm{s}}{\tau_B} \right)^{1/3} \mathrm{GeV}$$

$$100\ s \rightarrow \mathrm{TeV}$$

The high energy tail

$$T \simeq 7.8 \times 10^3 \times \left(\frac{1\ \mathrm{s}}{\tau_B} \right)^{1/3} \mathrm{GeV}$$

$$100\ s \rightarrow \mathrm{TeV}$$
$$15\ y \rightarrow 10\ \mathrm{GeV}$$

Can the KM3NeT be a PBH burst?



Can the KM3NeT be a PBH burst?





Can the KM3NeT be a PBH burst?

HAWC follow-up observations of the KM3NeT neutrino event KM3-230213A

ATel #17069; *Dezhi Huang (UMD) for the HAWC Collaboration*
*on **7 Mar 2025; 17:31 UT***
Credential Certification: Dezhi Huang (dezhih@umd.edu)

Subjects: Gamma Ray, TeV, Neutrinos

𝕏 Post

We report results from the High Altitude Water Cherenkov (HAWC) Gamma-ray Observatory on the extremely high-energy neutrino event KM3-230213A recently published by the KM3NeT Collaboration (Nature, 2024).

HAWC conducted a search for gamma-ray emission at the neutrino event position reported by KM3NeT over four different intervals. As the HAWC Observatory was undergoing maintenance during the KM3NeT detection, HAWC's follow-up observations began approximately 21 hours after the neutrino event. The event location first entered HAWC's field of view around February 14, 2023, at 00:50:00 UTC, with two complete transit covered between 21 and 69 hours after the KM3NeT detection. Subsequent searches were conducted one week, one month, and one year after the event, each with an average exposure of approximately 5 hours per transit as the event location passed through HAWC's field of view. Using a likelihood analysis and assuming a power-law spectral model with a fixed index of -2.7 and the fixed position reported by KM3NeT, no significant gamma-ray emission was detected in any of the intervals. We, therefore, derived the following 95% confidence-level upper limits on the gamma-ray flux (I_0) at 1 TeV:





Can the KM3NeT be a PBH burst?





Caveat: Spoiler alert! Pre burst monitors





One PBH event in KM3NeT would not be serendipitous

ARCA is optimized for the TeV range



One PBH event in KM3NeT would not be serendipitous

ARCA is optimized for the TeV range

KM3NeT is not fully assembled



One PBH event in KM3NeT would not be serendipitous

ARCA is optimized for the TeV range

KM3NeT is not fully assembled

One event in KM3NeT requires:

$$d \sim 4 \times 10^{-5}\ \mathrm{pc}$$



One PBH event in KM3NeT would not be serendipitous



ARCA is optimized for the TeV range

KM3NeT is not fully assembled

One event in KM3NeT requires:

$$d \sim 4 \times 10^{-5} \text{ pc}$$



Back to square one

Can neutrinos be the only candidate for observing a source?

Back to square one

Can neutrinos be the only candidate for observing a source?

Gamma rays can be:

Back to square one

Can neutrinos be the only candidate for observing a source?

Gamma rays can be:

Trapped or attenuated

Wang, arxiv:2310.15832
Meszaros, arxiv:1904.04226
Doi, arxiv:1909.02239
Ahlers, arxiv:1509.00805

Back to square one

Can neutrinos be the only candidate for observing a source?

Gamma rays can be:

Trapped or attenuated

Wang, arxiv:2310.15832
Meszaros, arxiv:1904.04226
Doi, arxiv:1909.02239
Ahlers, arxiv:1509.00805

Delayed

Hooper, arxiv:2502.09545

Back to square one

Can neutrinos be the only candidate for observing a source?

Gamma rays can be:

Trapped or attenuated

Wang, arxiv:2310.15832
Meszaros, arxiv:1904.04226
Doi, arxiv:1909.02239
Ahlers, arxiv:1509.00805

Delayed

Hooper, arxiv:2502.09545

Outside the field of view

Back to square one

Can neutrinos be the only candidate for observing a source?

Gamma rays can be:

Trapped or attenuated

Wang, arxiv:2310.15832
Meszaros, arxiv:1904.04226
Doi, arxiv:1909.02239
Ahlers, arxiv:1509.00805

Delayed

Hooper, arxiv:2502.09545

Outside the field of view

Observing the universe through the neutrino lens



Observing the universe through the neutrino lens





Neutrino telescope synergy



$\mathrm{RA_{PBH}} = 300^\circ$ (20h)

- IceCube – Gen2
- KM3NeT
- Baikal – GVD
- P – ONE
- TRIDENT

13th of February 2023

- 01 : 16 : 47 UTC
- 11 : 16 : 47 UTC



What did we learn today?



What did we learn today?



What did we learn today?



What did we learn today?



What did we learn today?



What did we learn today?